

DIVIDE TO CONQUER[®] THE DAY



DHIVY is a carbidopa/levodopa (CD/LD) tablet made to be broken—providing personalized and precise daily dosing for people with Parkinson's disease



Images do *not* represent the actual size of the tablet.

INDICATION

DHIVY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease-like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

DHIVY is contraindicated in patients

- Currently taking a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine, linezolid, and tranylcypromine) or have recently (within 2 weeks) taken a nonselective MAO inhibitor. Hypertension can occur if these drugs are used concurrently.
- With known hypersensitivity to any component of DHIVY.


Please see Important Safety Information throughout and accompanying full Prescribing Information.

People with Parkinson's disease are often prescribed CD/LD

CD/LD is the standard of care for motor symptoms associated with Parkinson's disease

Finding the right daily dose of CD/LD can be a challenge

- **Too much CD/LD** may cause dyskinesia (uncontrolled, involuntary muscle movement)
- **Not enough CD/LD** may cause OFF time (worsening of symptoms)



Your prescribed daily dose of CD/LD should help you manage your symptoms

Your doctor may recommend dividing your daily dose of CD/LD

Smaller, more frequent doses of oral CD/LD have been shown to help minimize dyskinesia and OFF time



Pill splitting can be imprecise

When you split a pill that isn't designed to be broken, you may be taking an inaccurate dose.



Pill splitting can be difficult

Pills are not easy to cut or break without crumbling.


Always talk to your doctor before adjusting your daily dose of CD/LD

DHIVY was designed with your unique needs in mind

2 doctors developed DHIVY using a patented technology

Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg

DHIVY is functionally scored,
so it's made to be broken



With deep grooves, DHIVY was designed to break evenly and cleanly

Firm enough **not to break accidentally**

Image does *not* represent the actual size of the tablet.

IMPORTANT SAFETY INFORMATION (continued)

Taking DHIVY may result in falling asleep while engaged in normal activities, even without warning and as late as 1 year or later after starting DHIVY. This may affect your ability to drive or operate machinery, resulting in accidents. Do not do anything that requires alertness until you know how DHIVY affects you. Tell your healthcare provider about any sleep medicines you take or if you have a sleep disorder, and discuss any episodes of drowsiness or sleepiness you experience while taking DHIVY.

Talk to your healthcare provider before you change the dose or stop taking DHIVY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle stiffness.

Please see Important Safety Information throughout and accompanying full Prescribing Information.

DHIVY provides daily dosing flexibility at your fingertips


Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg

DHIVY can be taken whole or in segments

Take DHIVY as prescribed by your doctor.

If you need a whole tablet of DHIVY:


Swallow the tablet whole. If you have trouble swallowing a whole tablet, you can snap DHIVY into segments.




If you need to snap DHIVY into segments:

Ensure you are holding the tablet in both hands, then gently push downward on each side of the tablet's groove.

Snapping a tablet into quarters



Snapping a tablet in half



Images do *not* represent the actual size of the tablet.

IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider if you have any heart conditions, especially if you have had a heart attack or irregular heartbeats. Your heart function should be monitored during the time period your DHIVY dose is first adjusted.

Please see Important Safety Information throughout and accompanying full Prescribing Information.

DHIVY is a CD/LD tablet designed to be divided

In one snappable tablet, DHIVY provides a range of segments

Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg

Divide your daily dose to fit your day



1 whole tablet
(25 mg/100 mg)



3/4 tablet
(18.75 mg/75 mg)



1/2 tablet
(12.5 mg/50 mg)



1/4 tablet
(6.25 mg/25 mg)

1/4 of DHIVY is the smallest segment available within a 25 mg/100 mg CD/LD tablet

Images represent the actual size of the tablet.

Talk to your doctor about dividing your daily CD/LD dose

IMPORTANT SAFETY INFORMATION (continued)

Treatment with DHIVY may cause hallucinations or abnormal thoughts and behaviors. Be sure to tell your healthcare provider if you experience excessive suspicion; seeing, hearing, or feeling things that are not real; confusion; agitation; aggressive behavior; and disorganized thinking.

Please see Important Safety Information throughout and accompanying full Prescribing Information.

DHIVY is a tool to help you take an accurate daily dose of CD/LD

Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg

Work closely with your doctor to create a daily dosing schedule



*For additional tips, download the
DHIVY Success Guide at DHIVY.com*

IMPORTANT SAFETY INFORMATION (continued)

Treatment with DHIVY may also cause intense urges to gamble, increased sexual urges, intense urges to spend money, binge eating, and other intense urges, as well as the inability to control those urges. Uncontrolled sudden movements of the face, arms, legs, or trunk may appear or get worse during treatment with DHIVY. This may mean your dosage of DHIVY or other Parkinson's medications needs to be adjusted.

Please see Important Safety Information throughout and accompanying full Prescribing Information.

Daily dosing reminders

Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg



Snap off only the amount you need



Place the remainder of the DHIVY tablet in a container protected from light and moisture



Take DHIVY with or without food



Store DHIVY at room temperature

Talk to your doctor before making any adjustments to your medications



IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider if you have ever had a peptic ulcer or glaucoma, as treatment with DHIVY may worsen these conditions. DHIVY also may cause abnormal blood or urine test results.

Some patients taking DHIVY may experience depression or suicidal thoughts. Tell your healthcare provider if you have thoughts of suicide.

The most common side effects that may appear with DHIVY include uncontrolled sudden movements and nausea.

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using DHIVY with certain other medicines, including MAO inhibitors, isoniazid, iron supplements, medicines for high blood pressure or those that increase or decrease dopamine levels, or metoclopramide, may cause serious side effects.

Notify your healthcare provider if you become pregnant or intend to become pregnant during DHIVY therapy or if you intend to breastfeed or are breastfeeding an infant.

Please see Important Safety Information throughout and accompanying full Prescribing Information.

The side effects of DHIVY are similar to other forms of CD/LD



The most common side effects associated with DHIVY:

- Dyskinesia
- Nausea

Talk to your doctor about any side effects you experience while taking DHIVY

IMPORTANT SAFETY INFORMATION

DHIVY is contraindicated in patients

- Currently taking a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine, linezolid, and tranylcypromine) or have recently (within 2 weeks) taken a nonselective MAO inhibitor. Hypertension can occur if these drugs are used concurrently.
- With known hypersensitivity to any component of DHIVY.

Taking DHIVY may result in falling asleep while engaged in normal activities, even without warning and as late as 1 year or later after starting DHIVY. This may affect your ability to drive or operate machinery, resulting in accidents. Do not do anything that requires alertness until you know how DHIVY affects you. Tell your healthcare provider about any sleep medicines you take or if you have a sleep disorder, and discuss any episodes of drowsiness or sleepiness you experience while taking DHIVY.

Talk to your healthcare provider before you change the dose or stop taking DHIVY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle stiffness.

Tell your healthcare provider if you have any heart conditions, especially if you have had a heart attack or irregular heartbeats. Your heart function should be monitored during the time period your DHIVY dose is first adjusted.

Treatment with DHIVY may cause hallucinations or abnormal thoughts and behaviors. Be sure to tell your healthcare provider if you experience excessive suspicion; seeing, hearing, or feeling things that are not real; confusion; agitation; aggressive behavior; and disorganized thinking.

Treatment with DHIVY may also cause intense urges to gamble, increased sexual urges, intense urges to spend money, binge eating, and other intense urges, as well as the inability to control those urges. Uncontrolled sudden movements of the face, arms, legs, or trunk may appear or get worse during treatment with DHIVY. This may mean your dosage of DHIVY or other Parkinson's medications needs to be adjusted.

Please see Important Safety Information throughout and accompanying full Prescribing Information.

There are various ways to save on a DHIVY prescription

See which savings option is the best for you at [DHIVY.com/savings-and-support/](https://www.dhivy.com/savings-and-support/)

Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg



Find more resources and support at [DHIVY.com](https://www.dhivy.com)

IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider if you have ever had a peptic ulcer or glaucoma, as treatment with DHIVY may worsen these conditions. DHIVY also may cause abnormal blood or urine test results.

Some patients taking DHIVY may experience depression or suicidal thoughts. Tell your healthcare provider if you have thoughts of suicide.

The most common side effects that may appear with DHIVY include uncontrolled sudden movements and nausea.

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using DHIVY with certain other medicines, including MAO inhibitors, isoniazid, iron supplements, medicines for high blood pressure or those that increase or decrease dopamine levels, or metoclopramide, may cause serious side effects.

Notify your healthcare provider if you become pregnant or intend to become pregnant during DHIVY therapy or if you intend to breastfeed or are breastfeeding an infant.

Please see Important Safety Information throughout and accompanying full Prescribing Information.

Take on the day with DHIVY

CD/LD designed with your unique needs in mind

Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg



Made to be broken—in one snappable tablet, DHIVY provides a range of segments



DHIVY provides personalized and accurate CD/LD daily dosing to fit your day



Work with your doctor to create a daily dosing schedule

See if you're eligible to save on DHIVY by visiting [DHIVY.com](https://www.dhivy.com)



INDICATION

DHIVY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease-like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

DHIVY is contraindicated in patients

- Currently taking a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine, linezolid, and tranylcypromine) or have recently (within 2 weeks) taken a nonselective MAO inhibitor. Hypertension can occur if these drugs are used concurrently.
- With known hypersensitivity to any component of DHIVY.

Please see Important Safety Information throughout and accompanying full Prescribing Information.

Avion
PHARMACEUTICALS

DHIVY and "Divide to Conquer" are registered trademarks of Avion Pharmaceuticals, LLC.
© 2023 Avion Pharmaceuticals, LLC
Alpharetta, GA 30005 | 1-888-612-8466 | 1905-v2